

comment on: " Even-odd behavior of conductance in monatomic sodium wires"

(February 1, 2008)

Based on numerical **LDA** calculation in their recent letter Sim et al. report an even-odd behavior of the conductance as a function of the number of atoms in monatomic sodium wires. [1] For a string of up to $N = 5$ sodium atoms with ends connected to leads with mirror symmetry, they found the conductance to have the value $\mathcal{G}_0 = 2e^2/h$ for odd N , while for even N it is smaller and sensitive to the wire structure. We here give a simple analytic proof of this result, and demonstrate that such an even-odd behavior exists in more general cases. We also argue that this parity effect holds for larger N as long as N is finite.

We model the wire as a string of N s-orbital (Na , Au , Ag , Cu) atoms [2] with its ends connected to leads, by the following tight-binding hamiltonian

$$H = \sum_{i=1}^N \epsilon_s a_{is}^+ a_{is} + \sum_{i=1}^{N-1} (t_i a_{is}^+ a_{i+1s} + H.c.) + \sum_{k,r=L,R} \epsilon_{kr} b_{kr}^+ b_{kr} + \sum_k (V_{ks}^L b_{kL}^+ a_{1s} + V_{ks}^R b_{kR}^+ a_{Ns} + H.c.), \quad (1)$$

where $a_{is}(b_{kr})$ is the annihilation operator of an s-orbital electron at atomic site i (lead r), and the other terms have their usual meaning. The Landauer-Büttiker conductance is determined by the transmission probability at the Fermi level, $\mathcal{G} = \mathcal{G}_0 \mathcal{T}(E_F)$. Considering the atomic sites to the right of site 1 as part of the right lead, the Keldysh formalism yields for the transmission probability the expression,

$$\mathcal{T}(E_F) = \frac{-2\Gamma^L |t_1|^2 \text{Im} G_{2R}^r}{(E_F - \epsilon_s - |t_1|^2 \text{Re} G_{2R}^r)^2 + (\Gamma_L - 2|t_1|^2 \text{Im} G_{2R}^r)^2/4}. \quad (2)$$

where

$$\begin{aligned} \Gamma^{L/R} &= \sum_k 2\pi |V_{ks}^{L/R}|^2 \delta(\epsilon - \epsilon_{kL/R}), \\ G_{iR}^r(E_F) &= [g_{is}^r(E_F)]^{-1} - |t_i|^2 G_{i+1,R}^r(E_F)]^{-1}, \quad i = 2, 3, \dots, N-1 \\ G_{NR}^r(E_F) &= [g_{NN}^r(E_F)]^{-1} + \frac{i}{2} \Gamma^R]^{-1} \\ g_{is}^r(E_F) &= (E_F - \epsilon_s + i0^+)^{-1}. \quad i = 1, 2, \dots, N \end{aligned}$$

At resonance ($E_F = \epsilon_s$) the real part of all retarded Green functions becomes zero and the first term in the denominator of Eq. (2) vanishes. Complete transmission through the dot lattice is then obtained if

$$\Gamma^L = -2|t_1|^2 \text{Im} G_{2R}^r, \quad (3)$$

where now

$$ImG_{2R}^r = -\frac{\Gamma^R}{2} \left| \frac{t_3 t_5 \cdots t_{N-2}}{t_2 t_4 \cdots t_{N-1}} \right|^2; \quad N \text{ odd} \quad (4)$$

$$ImG_{2R}^r = -\frac{2}{\Gamma^R} \left| \frac{t_3 t_5 \cdots t_{N-1}}{t_2 t_4 \cdots t_{N-2}} \right|^2; \quad N \text{ even}. \quad (5)$$

The condition thus becomes

$$\left| \frac{t_1 t_3 \cdots t_{N-2}}{t_2 t_4 \cdots t_{N-1}} \right|^2 = \frac{\Gamma^L}{\Gamma^R}; \quad N \text{ odd} \quad (6)$$

$$\left| \frac{t_1 t_3 \cdots t_{N-1}}{t_2 t_4 \cdots t_{N-2}} \right|^2 = \frac{\Gamma^L \Gamma^R}{4}; \quad N \text{ even}. \quad (7)$$

Equation (6) states that for a sample with an odd number of atoms and mirror symmetry ($\Gamma^L = \Gamma^R, t_1 = t_{N-1}, t_2 = t_{N-2}, \text{etc.}$) complete transmission is automatically satisfied, yielding $\mathcal{G} = 2e^2/h$. This is not the case when N is even, however, as is apparent from the structure of Eq. (7). For a wire with deformations perpendicular to its length the even-odd character of conductance is preserved since the normal deformation just changes the inter-site couplings symmetrically. One also finds that the conductance quantization for even N is more sensitive to the wire structure.

The above results are valid for any finite N . Also, they appear not to arise from the electron-electron interaction since our derivation ignores such coupling. If the latter is added, however, we expect them to hold as well since the on-site Coulomb interaction introduces a self-energy term Σ_{e-e} in the Green's function of each site. The influence of electron-electron interactions is then just to shift and split the resonance position [3] and Eq. (2) can also be formally used in their presence, with the formal replacement $g_{is}^r = (E_F - \epsilon_s - Re\Sigma_{e-e})^{-1}$ and the new resonance condition $\epsilon_s = E_F - Re\Sigma_{e-e}$.

This work was supported in part by a Catedra Presidencial en Ciencias and FONDECYT 1990425 (Chile), and NSF grant No. 53112-0810 of Hunan Normal University (China).

Z. Y. Zeng^{1,2} and F. Claro¹

¹Facultad de Física, Pontificia Universidad Católica de Chile, Casilla 306, Santiago 22, Chile

²Department of Physics, Hunan Normal University, Changsha 410081, China

PACS numbers: : 73.40.Cg, 73.40.Jn, 73.63.Rt

REFERENCES

- [1] H. -S. Sim, H. -W. Lee, and K. J. Chang, Phys. Rev. Lett. **87**, 096803 (2001).
- [2] A. Levy Yeyati, A. Martín-Rodero, and F. Flores, Phys. Rev. B **56**, 10369 (1997).
- [3] D. C. Langreth, Phys. Rev. **150**, 516 (1966).